



## SECTION 'C'

### Q.7 Multiple Choice Questions. ( MCQ )

1. A commercial method of grafting used in propagation of mango.
  - a. Inarch grafting
  - b. Bridge Cutting
  - c. Stone Grafting
  - d. None of these
2. Tissue culture method of propagation has been successfully and commercially used in
  - a. Banana
  - b. Ber
  - c. Sapota
  - d. All of these
3. Sexually propogated fruit crop is.
  - a. Papaya
  - b. Pineapple
  - c. Peach
  - d. Pear
4. Growth regulator used for rooting of cutting
  - a. GA
  - b. NAA
  - c. IBA
  - d. ABA
5. Parts of mother plants each of which is capable of producing roots in contact with soil while connect with mother plant is known as
  - a. Sucker
  - b. Runner
  - c. Bulb
  - d. Corm
6. Growth regulator used for breaking the seed dormancy
  - a. GA
  - b. ABA
  - c. IBA
  - d. All of these
7. A mass of parenchyma cells that develop from and around the wounded plant tissue is
  - a. Stolens
  - b. Tubers
  - c. offsetts
  - d. Callus
8. A regeneration of adventitious roots from a shoot when it is still attached with the mother plant.
  - a. Budding
  - b. Layering
  - c. Grafting
  - d. Cutting
9. Root stock used for budding of mandarin orange
  - a. Khirni
  - b. Rayon
  - c. Wood apple
  - d. Jambheri
10. New plant resulting from union of male and female gametes during fertilization is called.
  - a. Seed
  - b. Embryo
  - c. Seed coat
  - d. None of these

11. Pineapple is commercially propagated by
- a. Seed
  - b. Suckers
  - c. Rhizome
  - d. Eyebuds
12. The seeds are arranged in alternate layers of moist sand or soil and stored at 32<sup>0</sup> F - 45<sup>0</sup> F for 1-4 month .
- a. Scarification
  - b. Scalding
  - c. Stratification
  - d. None of these

==x0x==